

$[SU(3) \times SU(2) \times U(1)]^2$ and Strong Unification

Chun Liu

Institute of Theoretical Physics, Chinese Academy of Sciences,

P. O. Box 2735, Beijing 100080, China

Abstract

A supersymmetric model with gauge symmetry $G_1 \times G_2$, where $G_i = SU(3)_i \times SU(2)_i \times U(1)_i$, is constructed within the framework of gauge mediated supersymmetry breaking. At the energy scale $\sim (10 - 100)$ TeV where the gauge symmetry breaks down to the Standard Model (SM), G_1 is strong and G_2 is weak. The observed gauge coupling constant unification of the SM is attributed to that of G_2 . The messenger fields and Higgs fields just satisfy the condition that makes G_2 a realization of strong unification. The SM gauginos are predicted to be generally heavier than the sleptons and squarks.

Keywords: gauge interaction, supersymmetry.

PACS numbers: 12.60.-i, 12.60.Cn, 12.60.Jv

Extensions of the SM aim at understanding new experimental results or unsolved theoretical problems. The most popular approach is the grand unification theories (GUTs) [1], such as the $SU(5)$ GUT. There are indirect experimental evidences for GUTs from LEP and neutrino physics. To make GUTs viable, supersymmetry (SUSY) [2,3] is a must. One of the novel idea towards GUT is the so-called strong unification [4,5]. In the strong GUT, the SM gauge coupling constants just reach their common Landau pole at the unification energy scale.

Strong GUT is interesting not only due to its novelty, but also because of its usefulness. There is a discrepancy between the measured value of the QCD strong coupling constant at M_Z , which is $\alpha_s^{\text{exp}}(M_Z) \simeq 0.1172 \pm 0.002$ [6], and that predicted by the minimal SUSY SM (MSSM) $\alpha_s^{\text{MSSM}}(M_Z) \simeq 0.126$. The discrepancy is reduced if extra matters are added into the MSSM. To keep the unification, the additional states should be in complete representation of GUT gauge groups. To the two-loop level, it has been shown [5], for example, that $\alpha_s(M_Z) \simeq 0.1163$ if there are additional six multiplets in $\mathbf{5} + \bar{\mathbf{5}}$ under $SU(5)$ with masses of ~ 214 TeV. However, the model would be artificial if these additional matters are naively added.

We will illustrate that a SUSY model with the gauge symmetry $SU(3)_1 \times SU(2)_1 \times U(1)_1 \times SU(3)_2 \times SU(2)_2 \times U(1)_2$ can be a nontrivial realization of the strong GUT. There are multiple motivations to consider such an extension of the SM [7–9]. In Ref. [9], such kind of models were proposed as GUT generalization of the SUSY top-color [10]. They provide a solution to the SUSY flavor changing neutral current problem. However, the gauge coupling constant behavior was rather bad at high energies because of the introduction of too many extra matter fields which made the gauge interactions to be too much strong. This situation brings us to further think of their connection with the idea of the strong GUT. In this paper, after naturally modifying the Higgs and messenger contents of the model, we note that the extra matters additional to the MSSM can make the SM-like gauge interaction $SU(3)_2 \times SU(2)_2 \times U(1)_2$ a strong GUT.

We consider a SUSY theory with the gauge group $G_1 \times G_2$ in the framework of gauge

mediated SUSY breaking (GMSB) [11], where $G_i = SU(3)_i \times SU(2)_i \times U(1)_i$ ($i = 1, 2$). The three coupling constants of G_1 are large, and those of G_2 are small at the TeV scale. The three generations of matter carry nontrivial quantum numbers of G_2 only. These numbers are assigned in the same way as they are under the SM gauge group. One gauge singlet chiral superfield X is introduced for SUSY breaking,

$$\langle X_s \rangle \neq 0, \quad \langle F_X \rangle \neq 0, \quad (1)$$

with X_s and F_X being the scalar and auxiliary components of X . The vacuum expectation values are taken to be real.

For the SUSY breaking messengers and gauge symmetry breaking Higgs', it is easy to consider them through imaging global $SU(5)_i$ symmetry into which G_i is embedded. The messengers with their quantum numbers under $SU(5)_1 \times SU(5)_2$ are

$$\begin{aligned} T_1(5, 1), \quad \bar{T}_1(\bar{5}, 1), \\ T_2(1, 5), \quad \bar{T}_2(1, \bar{5}) \end{aligned} \quad (2)$$

The relevant superpotential is

$$\mathcal{W}_1 = c_1 X T_1 \bar{T}_1 + c_2 X T_2 \bar{T}_2, \quad (3)$$

where c_1 and c_2 are coupling constants of order one. The fields T_i and \bar{T}_i are massive at tree level. Their fermionic components compose a Dirac fermion with mass $c_i \langle X_s \rangle$, while the scalar components have a squared-mass matrix

$$(T_{is}^* \quad \bar{T}_{is}) \begin{pmatrix} c_i^2 \langle X_s \rangle^2 & c_i \langle F_X \rangle \\ c_i \langle F_X \rangle & c_i^2 \langle X_s \rangle^2 \end{pmatrix} \begin{pmatrix} T_{is} \\ \bar{T}_{is}^* \end{pmatrix}. \quad (4)$$

The mass eigenstates and squared-mass eigenvalues are

$$\begin{aligned} \frac{1}{\sqrt{2}}(T_{is} + \bar{T}_{is}^*) \quad \text{with} \quad m_{i1}^2 = c_i^2 \langle X_s \rangle^2 + c_i \langle F_X \rangle, \\ \frac{1}{\sqrt{2}}(T_{is} - \bar{T}_{is}^*) \quad \text{with} \quad m_{i2}^2 = c_i^2 \langle X_s \rangle^2 - c_i \langle F_X \rangle. \end{aligned} \quad (5)$$

It is assumed that $c_i \langle F_X \rangle < c_i^2 \langle X_s \rangle^2$. Because $\langle F_X \rangle \neq 0$, SUSY breaking occurs in the fields T_i 's and \bar{T}_i 's at tree-level. G_1 and G_2 sectors get to be soft SUSY breaking via the

messengers at loop level. Because G_2 is weak at TeV scale, its SUSY breaking effects can be calculated perturbatively, for example, G_2 gaugino soft masses are

$$M_{\lambda_r} \simeq \frac{\alpha'_r}{4\pi} \frac{\langle F_X \rangle}{\langle X_s \rangle}, \quad (6)$$

where $\alpha'_r = g'_r/4\pi$ with g'_r being the gauge coupling constants of G_2 . And $r = 1, 2, 3$ corresponding to the groups $U(1)$, $SU(2)$, and $SU(3)$, respectively. However, G_1 is strong, we can only estimate its gaugino masses

$$M_{\lambda_r} \simeq \frac{\langle F_X \rangle}{\langle X_s \rangle}. \quad (7)$$

Numerically the messenger masses are about $(10 - 100)$ TeV.

A pair of Higgs $\Phi_1(5, \bar{5})$ and $\Phi_2(\bar{5}, 5)$ breaks the $G_1 \times G_2$ gauge symmetry down to that of the SM. One gauge singlet superfield Y is introduced for the gauge symmetry breaking. The superpotential of them is written as follows,

$$\mathcal{W}_2 = c' Y [\text{Tr}(\Phi_1 \Phi_2) - \mu'^2], \quad (8)$$

where the trace is taken with regard to both $SU(3)_1 \times SU(3)_2$ and $SU(2)_1 \times SU(2)_2$. μ' is the energy scale relevant to the gauge symmetry breaking, and c' is the coupling constant. The Higgs fields get soft masses like that given in Eq. (7). However, the above superpotential is not enough to guarantee all the Φ_i fermion components to be massive. Their masses are nonvanishing when a superfield A which is in adjoint representation of $SU(5)_1$ is introduced with the following superpotential [8],

$$\mathcal{W}'_2 = c'_2 \text{Tr}(\Phi_2 A \Phi_1) \quad (9)$$

with c'_2 being the coupling constant. The details of the gauge symmetry breaking go the same way as that in Ref. [9] (its Eqs. (10-17)). The VEVs of the Φ_i 's are given as

$$\langle \Phi_{1s} \rangle = \langle \Phi_{2s} \rangle = v I_3 \otimes I_2, \quad (10)$$

where I_3 and I_2 are the unit matrices in the space of $SU(3)_1 \times SU(3)_2$ and $SU(2)_1 \times SU(2)_2$, respectively. The coupling constants of the SM $SU(3)_c \times SU(2)_L \times U(1)_Y$ are

$$\frac{1}{g_s^2} = \frac{1}{g_3^2} + \frac{1}{g_3'^2}, \quad \frac{1}{g^2} = \frac{1}{g_2^2} + \frac{1}{g_2'^2}, \quad \frac{1}{g'^2} = \frac{1}{g_1^2} + \frac{1}{g_1'^2}. \quad (11)$$

Numerically, the gauge symmetry breaking scale v is about $(10 - 100)$ TeV.

Electroweak symmetry breaking is achieved via a pair of Higgs superfields H_u and H_d which are nontrivial only under G_2 [9].

Around $10 - 100$ TeV, there are many matter fields which will run the gauge coupling constants to be large at high energies. The matter fields introduced additional to MSSM are complete $SU(5)$ multiplets. Therefore, the unification scale 3×10^{16} GeV is still the same as that of the MSSM, but the values of the coupling constants are significantly different. This model is a candidate of strong GUT.

Below the scale v , the $G_1 \times G_2$ breaks spontaneously down to the MSSM. From Eq. (11), it is easy to see that the gauge coupling constants of MSSM are almost fully determined by that of G_2 , because $g_i \gg g_i'$. Therefore the observed unification of MSSM is attributed to the unification of G_2 .

Above $(10 - 100)$ TeV scale, the theory is $G_1 \times G_2$. As far as the G_2 sector is concerned, the new matter fields in addition to the MSSM are the messengers T_1 and \bar{T}_1 , and Higgs fields Φ_1 and Φ_2 . The messenger fields compose one $\mathbf{5} + \bar{\mathbf{5}}$ multiplet with a mass $c\langle X_s \rangle$ and the Higgs' contribute five $\mathbf{5} + \bar{\mathbf{5}}$ multiplets with masses $c'v$ as well as c'_2v . We have the freedom to adjust all the masses of these six $\mathbf{5} + \bar{\mathbf{5}}$ multiplets to be about 214 TeV. As has been shown in Ref. [5], the gauge couplings reach their common Landau pole at the GUT scale $\sim 3 \times 10^{16}$ GeV. Namely in this case, G_2 is a realization of the strong GUT.

Some remarks are necessary. (1) The perturbative calculation in Ref. [5] was not reliable around the GUT scale because of the large coupling constants. But around 100 TeV where the perturbative domain lies, its reliability was under control. It is in the latter low energy region where we have made use of Ref. [5]. (2) On the other hand, G_1 sector is also expected to be a GUT. $(10 - 100)$ TeV is already its non-perturbative region, we have no reliable method yet to make detailed analysis. (3) The unification simply means that the gauge coupling constants are equal at certain scales. We have not introduced any unified gauge

group. Such a model does not have proton decays, and does not suffer from the doublet-triplet splitting problem. (4) It should be noted that only is G_2 SM-like, can the breaking $G_1 \times G_2 \rightarrow \text{SM}$ at $(10 - 100)$ TeV occur. Any breaking of $SU(5) \times SU(5) \rightarrow SU(5)$ [12] would have occurred above 3×10^{16} GeV. (5) Some of the matter contents of G_2 , such as the third generation can be moved into G_1 . Due to GMSB, the superpartners in this sector are very heavy $\sim 10 - 100$ TeV. They decouple at $(1 - 10)$ TeV energy scale. At this low energy scale the fermions, on the other hand, can form condensates due to the strong gauge interactions. Dynamical fermion masses might be generated [10]. In order to keep the strong GUT, it is possible to either introduce one more $\mathbf{5} + \bar{\mathbf{5}}$ multiplet of G_2 , which may play a role of SUSY breaking messengers [9], or lower the SUSY breaking and messenger scales to be around 10 TeV. These possibilities should be studied further and are beyond the scope of this work. (6) If the $SU(3)_1$ interaction is switched off, the model is a kind of top-flavor models [13].

This model has interesting phenomenology. Besides the new gauge bosons, gauginos and Higgs particles with masses around $(10 - 100)$ TeV, the SM gaugino masses are predicted to be as heavy as $\sim 100 \text{ GeV} - 1 \text{ TeV}$. Let us analyze the gaugino spectrum in more detail. The full gaugino masses have two origins: SUSY breaking (soft masses) and spontaneous gauge symmetry breaking. It has been obtained in Ref. [9] that the relevant mass matrix in the basis of λ_r , λ'_r and the higgsino $(\psi_1 - \psi_2)/\sqrt{2}$ is

$$M_r = \begin{pmatrix} M_{\lambda_r} & 0 & \sqrt{2}g_r v \\ 0 & M_{\lambda'_r} & \sqrt{2}g'_r v \\ \sqrt{2}g_r v & \sqrt{2}g'_r v & 0 \end{pmatrix}, \quad (12)$$

where ψ_1 and ψ_2 stand for the fermion components of Φ_1 and Φ_2 , respectively. Numerically at the scale $v \sim (10 - 100)$ TeV, $g'_r \sim 0.1$, $g_r \sim 1$. The mass matrix determines two heavy states with masses $\sim M_{\lambda_r} \sim g_r v \sim (10 - 100)$ TeV, and one lighter state $\sim (g'_r v)^2/M_{\lambda_r} \sim 100 \text{ GeV} - 1 \text{ TeV}$. This lighter state is a mixture of the G_2 gaugino with the higgsino. It is regarded as the MSSM gaugino in this model. On the other hand, the soft masses of the three generation matters are about 100 GeV. Therefore in this model the SM gauginos

are generally heavier than the sleptons and squarks. Such a mass pattern can be tested in future colliders.

ACKNOWLEDGMENTS

The author acknowledges support from the National Natural Science Foundation of China.

REFERENCES

- [1] J.C. Pati and A. Salam, Phys. Rev. D10 (1974)275;
H. Georgi and S.L. Glashow, Phys. Rev. Lett. 32 (1974) 438.
- [2] J. Wess and B. Zumino, Nucl. Phys. B70 (1974) 39;
Y. Gol'fand and E. Likhtman, JETP Lett. 13 (1971) 323;
D.V. Volkov and V. Akulov, Phys. Lett. B46 (1973) 109.
- [3] E. Witten, Nucl. Phys. B188 (1981) 513.
- [4] L. Maiani, G. Parisi and R. Petronzio, Nucl. Phys. B136 (1978) 115;
N. Cabibbo and G.R. Farrar, Phys. Lett. 110B (1982) 107;
B. Brahmachari, U. Sarkar and K. Sridhar, Mod. Phys. Lett. A8 (1993) 3349;
R. Hempfling, Phys. Lett. B351 (1995) 206;
K.S. Babu and J.C. Pati, Phys. Lett. B384 (1996) 140;
C. Kolda, J. March-Russell, Phys. Rev. D55 (1997) 4252;
For a review, see V.A. Rubakov and S.V. Troitsky, hep-ph/0001213.
- [5] D. Ghilencea, M. Lanzafora and G.G. Ross, Phys. Lett. B 415 (1997) 253.
- [6] K. Hagiwara et al., Phys. Rev. D66 (2002) 010001.
- [7] N. Weiner, hep-ph/0106097;
N. Arkani-Hamed, A. G. Cohen, H. Georgi, hep-th/0108089;
E. Witten, hep-ph/0201018;
L.V. Laperashvili and H.B. Nielsen, Mod. Phys. Lett. A14 (1999) 2797;
Y. Kawamura, Prog. Theor. Phys. 105 (2001) 999.
- [8] H.-C. Cheng, D.E. Kaplan, M. Schmaltz and W. Skiba, Phys. Lett B515 (2001) 395.
- [9] J.G. Körner and C. Liu, Mod. Phys. Lett. A18 (2003) 967.
- [10] C. Liu, Phys. Rev. D61 (2000) 115001.

- [11] M. Dine and W. Fischler, Phys. Lett. B110 (1982) 227;
L. Alvarez-Gaumé, M. Claudson and M. Wise, Nucl. Phys. B207 (1982) 96;
C.R. Nappi and B.A. Ovrut, Phys. Lett. B113 (1982) 175. M. Dine and A.E. Nelson,
Phys. Rev. D48 (1993) 1277;
M. Dine, A.E. Nelson and Y. Shirman, Phys. Rev. D51 (1995) 1362;
M. Dine, A.E. Nelson, Y. Nir and Y. Shirman, Phys. Rev. D53 (1996) 2658.
- [12] For a recent study, see M. Dine, Y. Nir and Y. Shadmi, Phys. Rev. D66 (2002) 115001.
- [13] X. Li and E. Ma, Phys. Rev. Lett. 47 (1981) 1788.